# Understanding the basics of ALGS

This guide can help you understand Alagille (pronounced *a-luh-jeel*) syndrome, also known as ALGS, and a treatment for pruritus (itch), with tips for talking to doctors if you're not sure what to ask.

## ASK YOUR CHILD'S DOCTOR

**How is ALGS affecting my child's liver?**

**Are any tests needed to get a fuller picture of the impact of ALGS?**

**Do we need to visit other doctors who specialize in parts of the body affected by ALGS?**

### What is BYLVAY (odevixibat)?

BYLVAY is a prescription medicine for the treatment of itching in patients 12 months of age and older with Alagille syndrome (ALGS).

### What are the possible side effects of BYLVAY?

**BYLVAY can cause serious side effects, including:**

- You should not use BYLVAY if you have or have had an advanced stage of liver disease. Talk with your healthcare provider if you have any liver conditions.
- Abnormal liver tests have been observed with use of BYLVAY. Liver tests should be obtained before starting and periodically during therapy to monitor for liver injury. Immediately report to your healthcare provider any signs or symptoms of liver injury, such as nausea, vomiting, skin or the whites of eyes turn yellow, dark or brown urine, pain on the right side of the abdomen, or loss of appetite.

**Please see additional Important Safety Information throughout, and the full Prescribing Information and Instructions for Use for BYLVAY.**

Bylvay® (odevixibat)
200 | 400 | 600 | 1200 mcg capsules

# Talking to your doctor about treatment for itch from ALGS

## Ask your child's doctor about different ways to manage ALGS



**Nutritional supplements**



**Medications to treat itch**



**Surgery, liver transplant**

## BYLVAY is an FDA-approved medicine to treat pruritus (itch) in patients with ALGS as young as 12 months old

**Use the questions below as a starting point for talking to your doctor about whether BYLVAY is right for your child.**

### Could BYLVAY help my child with itch from ALGS?

### How is BYLVAY taken?

### How does BYLVAY work?

### What can we expect when starting BYLVAY?

### Can BYLVAY help with any level of itch?

### How soon can results be seen when starting BYLVAY?

### What side effects can occur when taking BYLVAY?

### What support is available for people starting BYLVAY?


### What are the possible side effects of BYLVAY? (cont'd)

**BYLVAY can cause serious side effects, including:**

- Notify your healthcare provider if you experience new onset or worsening of diarrhea.
- BYLVAY may decrease the absorption of certain vitamins, which include vitamins A, D, E, and K (FSV or fat-soluble vitamins). Your blood levels of these vitamins should be measured before starting and periodically during treatment to assess for FSV deficiency. You may bleed more easily, may bleed longer, or experience a bone fracture. Call your healthcare provider if you have any signs or symptoms of bleeding.

**Please see additional Important Safety Information throughout, and the full Prescribing Information and Instructions for Use for BYLVAY.**

Bylvay® (odevixibat)
200 | 400 | 600 | 1200 mcg capsules




# Bring this discussion guide to your next appointment to help you have a full conversation with your child's doctor about itch with ALGS

All images are actor portrayals.

## Talk to your doctor about how ALGS itch impacts your child

_____

_____

_____

## What are the possible side effects of BYLVAY? (cont'd)

**The most common side effects of BYLVAY include:**

- The most common side effects of BYLVAY treatment in patients with ALGS are diarrhea, abdominal pain, abnormal bleeding, such as bruising, and decreased weight.
- Do not swallow the 200 mcg or 600 mcg capsule(s) containing Oral Pellets whole. These are intended to be opened and the contents mixed into soft food.
- For patients taking bile acid binding resins, take BYLVAY at least 4 hours before or 4 hours after taking a bile acid binding resin.
- There is a pregnancy safety study that collects outcome data in women taking BYLVAY during pregnancy. Pregnant women exposed to BYLVAY should report if they have used BYLVAY during pregnancy by calling 1-855-463-5127 and contact your doctor immediately.

**You are encouraged to report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. You may also report side effects to Ipsen Biopharmaceuticals, Inc. at 1-855-463-5127.**

**Please see full Prescribing Information and Instructions for Use for BYLVAY.**

Bylvay® (odevixibat)
200 | 400 | 600 | 1200 mcg capsules

IPSEN

BYLVAY is a registered trademark of Albireo, Inc, an Ipsen company.
©2026 Ipsen Biopharmaceuticals, Inc.
All rights reserved. 08/2026 BYL-US-001712 v5.0.